

## Konventionelle und neue Parameter zur Abklärung eines Vitamin B12-Mangels

### Referenzen

1. Stabler SP. Clinical practice. Vitamin B12 deficiency. *N Engl J Med.* 2013;368(2):149-60.
2. Carmel R. Subclinical cobalamin deficiency. *Curr Opin Gastroenterol.* 2012;28(2):151-8.
3. Ebbing M, Bonna KH, Nygard O, Arnesen E, Ueland PM, Nordrehaug JE, et al. Cancer incidence and mortality after treatment with folic acid and vitamin B12. *JAMA.* 2009;302(19):2119-26.
4. Koenig V, Stanga Z, Zerlauth M, Bernasconi L, Risch M, Huber A, et al. Prevalence of vitamin B12 depletion and deficiency in Liechtenstein. *Public Health Nutr.* 2012:1-7.
5. Morkbak AL, Hvas AM, Milman N, Nexø E. Holotranscobalamin remains unchanged during pregnancy. Longitudinal changes of cobalamins and their binding proteins during pregnancy and postpartum. *Haematologica.* 2007;92(12):1711-2.
6. Multiple macrocytosis. *N Engl J Med.* 1967;277(11):595-6.
7. Foroozan P, Trier JS. Mucosa of the small intestine in pernicious anemia. *N Engl J Med.* 1967;277(11):553-9.
8. Risch C, Medina P, Nydegger UE, Bahador Z, Brinkmann T, Von Landenberg P, et al. The relationship of leukocyte anisocytosis to holotranscobalamin, a marker of cobalamin deficiency. *Int J Lab Hematol.* 2012;34(2):192-200.
9. Carmel R, Green R, Jacobsen DW, Qian GD. Neutrophil nuclear segmentation in mild cobalamin deficiency: relation to metabolic tests of cobalamin status and observations on ethnic differences in neutrophil segmentation. *Am J Clin Pathol.* 1996;106(1):57-63.
10. Oosterhuis WP, Niessen RW, Bossuyt PM, Sanders GT, Sturk A. Diagnostic value of the mean corpuscular volume in the detection of vitamin B12 deficiency. *Scand J Clin Lab Invest.* 2000;60(1):9-18.
11. Saxena S, Weiner JM, Carmel R. Red blood cell distribution width in untreated pernicious anemia. *Am J Clin Pathol.* 1988;89(5):660-3.

12. Carmel R, Agrawal YP. Failures of cobalamin assays in pernicious anemia. *N Engl J Med.* 2012;367(4):385-6.
13. Valente E, Scott JM, Ueland PM, Cunningham C, Casey M, Molloy AM. Diagnostic accuracy of holotranscobalamin, methylmalonic acid, serum cobalamin, and other indicators of tissue vitamin B(1)(2) status in the elderly. *Clin Chem.* 2011;57(6):856-63.
14. Nexo E, Hoffmann-Lucke E. Holotranscobalamin, a marker of vitamin B-12 status: analytical aspects and clinical utility. *Am J Clin Nutr.* 2011;94(1):359S-65S.
15. Herrmann W, Obeid R. Utility and limitations of biochemical markers of vitamin B12 deficiency. *Eur J Clin Invest.* 2013;43(3):231-7.
16. Carmel R. Holotranscobalamin: not ready for prime time. *Clin Chem.* 2012;58(3):643-5; author reply 5-6.
17. Hvas AM, Morkbak AL, Hardlei TF, Nexo E. The vitamin B12 absorption test, CobaSorb, identifies patients not requiring vitamin B12 injection therapy. *Scand J Clin Lab Invest.* 2011;71(5):432-8.
18. Carmel R. Biomarkers of cobalamin (vitamin B-12) status in the epidemiologic setting: a critical overview of context, applications, and performance characteristics of cobalamin, methylmalonic acid, and holotranscobalamin II. *Am J Clin Nutr.* 2011;94(1):348S-58S.
19. Fedosov SN. Metabolic signs of vitamin B(12) deficiency in humans: computational model and its implications for diagnostics. *Metabolism.* 2010;59(8):1124-38.