

Altershypogonadismus

Tilmann Möltgen, Andreas R. Huber, Kantonsspital Aarau

Referenzen

1. Kaufmann JM, Vermeulen A. The decline of androgen levels in elderly men and its clinical and therapeutic implications. *Endocr Rev* 2005 Oct;26(6):833-76
2. Wu FC, Tajar A, Pye SR, et al. Hypothalamic-pituitary-testicular axis disruptions in older men are differentially linked to age and modifiable risk factors: the European Male Aging Study. *J Clin Endocrinol Metab* 2008 Jul;93(7):2737-45
3. Erectile dysfunction and cardiovascular disease. Wespes E, Schulman CC. *Arch Esp Urol*. 2010 Oct;63(8):649-54.
4. Morley JE, Charlton E, Patrick P, et al. Validation of a screening questionnaire for androgen deficiency in aging males. *Metabolism* 2000 Sep;49(9):1239-42
5. Hall SA, Esche GR, Araujo AB, et al. Correlates of low testosterone and symptomatic androgen deficiency in a population-based sample. *J Clin Endocrinol Metab* 2008 Oct;93(10):3870-7
6. Wu FC, Tajar A, Beynon JM, et al. Identification of late-onset hypogonadism in middle aged and elderly men. *N Engl J Med* 2010 Jul 8;363(2):123-35
7. Veldhuis JD. Elements in the pathophysiology of diminished growth hormone (GH) secretion in aging humans. *Endocrine* 7:41;1997
8. **Abbildung A.** Herklotz R, Jermini F, Risch L, Huber AR. The aging male in Laboratory. *Labolife* 2002 6: Nov: 7-15.
9. **Abbildung B.** Hasler J, Herklotz R, Luppa PB, Diver MJ, Thevis M, Metzger J, Savoca R, Jermini F, Huber AR. Impact of recent biochemical findings on the determination of free and bioavailable testosterone: evaluation and proposal for clinical use. *J Lab Med* 2006 Jul;30(6):492-505